INDEX

ACOSTA, R. See Kemmerer, A. R., 527.

ALBANESE, A. A., S. E. SNYDERMAN, M. LEIN, E. M. SMETAK AND B. VESTAL. The biological value of corn and wheat proteins in the male infant, with a note on the utilization of D-tryptophan, 215.

ALLEN, D. I. See Swain, H. L., 63.

ALPER, C. See Chow, B. F., 319.

AMBROSE, A. M., AND F. DREDS. The value of rutin and quercetin in scurvy, 305.

AMERICAN INSTITUTE OF NUTRITION
Committee on Nomenclature, report of, 399.
— on Pathological Registries, report of, 398.
Editor, Journal of Nutrition, election of, 396.
— — — — report for 1948, 397.
Election of officers, 1949–50, 399.
New members elected, 400.
Proceedings, 13th annual meeting, 1949, 395.

Amino acid(s) and peptides, urinary excretion by dogs fed protein hydrolysates or amino acids, 155.
— — content, essential, of several vegetables, 527.
— — determination of some essential, in several uncooked and cooked Mexican foodstuffs, 293.
— — excretion by rats, and their growth, effect of fat in diet on, 257.
— — or protein hydrolysates, urinary excretion of amino acids and peptides by dogs fed, 155.
— — purified, as source of nitrogen for growing rat, 177.

Army (U. S.), simultaneous surveys of food consumption in various camps of, 51.

Artificial enrichment of white rice as solution to endemic beriberi. I. Preliminary report of field trials, 443.

ASENJO, C. F. See Goyco, J. A., 517.

Assay, microbiological, of "folic acid" as applied to milk, 25.

Award, B complex (see Mead Johnson and Co.).
— Borden, in nutrition (see Borden award).
— Osborne and Mendel (see Osborne and Mendel.)

BARKI, V. H. See Couch, J. R., 105.

BARNES, J. R. See Bratzler, J. W., 41.

BAUER, A. C. See Smith, S. E., 87.

B complex award (see Mead Johnson and Co.).

Beef, nutritive properties of protein in different cuts of, 381.

Beriberi (endemic), artificial enrichment of white rice as solution to: I. Preliminary report of field trials, 443.

Biological value(s) of corn and wheat proteins in male infant, with note on utilization of D-tryptophan, 215.
— — (relative to egg) and replacement values of some cereal proteins in human subjects, 405.

Biotin deficiency in mouse, production of, 73.

Birth, relation of maternal nutrition to condition of infant at: study of siblings, 453.

BLANDAU, R. J., H. KAUNITZ AND C. A. SLANETZ. Ovulation, fertilization, and transport of ova in old, vitamin E deficient rats, 97.

Blood concentrations, thiamine and pyruvic acid. Vitamin studies in middle-aged and old individuals. III., 353.
— of dairy cows, concentrations of various constituents in, during stages of terminal gestation and initial lactation. I. Effect of prepartal diet on serum tocopherols, 503.

Borden award in nutrition, presentation of, 1949, 402.

BOROUGHS, E. D. See Johnston, F. A., 479.

Boys (see Children).

BRATZLER, J. W., J. R. BARNES AND R. W. SWIFT. The effects of thiouracil on metabolism, 41.

BROWN, C. F. See Deuel, H. J., Jr., 361.

BURKE, B. S., S. S. STEVENSON, J. WORCESTER AND H. C. STUART. Nutrition studies during pregnancy. V. Relation of maternal nutrition to condition of infant at birth: study of siblings, 453.

CALBERT, C. E. See Deuel, H. J., Jr., 369.

Calcium metabolism (see Metabolism, calcium).

CALDWELL, A. B. See Taylor, C. M., 1.

CALVERLY, C. E., AND C. KENNEDY. The effect of fat, on calcium and phosphorus metabolism in normal growing rats under a normal dietary regime, 165.

CALVO, J. See Massieu, G. H., 293.

Carious lesions in cotton rat molars. III. Effect of particle size and consistency of purified ration on incidence and type of carious lesions, 275.

Casein or lactalbumin hydrolysates, effect of retention of nitrogen in, on regeneration of plasma proteins of protein-depleted dogs, 319.

Cereal proteins (some) in human subjects, relative to egg biological values and replacement values of, 405.

Chicks, influence of high levels of fat with suboptimum levels of riboflavin on growth of, 247.

CHIEFFI, M. See Kirk, E., 353.

Children 9 to 11 years of age, energy expenditure of: (1) sitting listening to the radio (phonograph), (2) sitting singing, and (3) standing singing. 1.

Choline requirement of baby pig, 195.

CHOW, B. F., C. ALPER AND S. DEBIASE. The effect of retention of nitrogen in casein or lactalbumin hydrolysates on the regeneration of plasma proteins of protein-depleted dogs, 319.
CLARK, L. C., H. RYER AND J. R. MURLIN. Relative to egg biological values and replacement values of some cereal proteins in human subjects, 405.
Climate, subarctic, food consumption of soldiers in (Fort Churchill, Manitoba, Canada, 1947–48), 63.
Cod liver, nutritive value of, 489.
— roe, nutritive value of, 489.
Consistency of purified ration and particle size, effect on incidence and type of carious lesions in cotton rat molars, 275.
CONSOLAZIO, F. C. See Swain, H. L., 63.
Corn and wheat proteins, biological value in male infant, with note on utilization of D-tryptophan, 215.
Cotton rat molars, carious lesions in. III., 275.
Cottonseed and rapeseed oils, digestibility in human subjects, 369
COUCH, J R., V. H. BARKI, M. L. SUNDE, W. W. CRAVENS AND C. A. ELVEHJEM. The utilization of lactose by the mature fowl, 105.
Cows, dairy, concentrations of various constituents in blood of, during stages of terminal gestation and initial lactation. I., 503.
CRAVENS, W. W. See Couch, J. R., 105.
CRAVIOTO, R. O. See Massieu, G. H., 293.

DAIRY cows (see Cows, dairy).
DEBIASE, S. See Chow, B. F., 319.
DEEDS, F. See Ambrose, A. M., 305.
DE LEON, J. F. See Salcedo, J., Jr., 443.
DEUEL, H. J., JR., S. M. GREENBERG, E. E. STRAUB, T. FUKUI, C. M. GOODING AND C. F. BROWN. Studies on the comparative nutritive value of fats. XI. On the possible growth-promoting activity of Δ^{12}-octadecenoic acid, 361.
— R. M. JOHNSON, C. E. CALBERT, J. GARDNER AND B. THOMAS. Studies on the comparative nutritive value of fats. XII. The digestibility of rapeseed and cottonseed oils in human subjects, 369.
Diet(s) of rats, effect of fat in, on growth and excretion of amino acids, 257.
— prepartal, effect on serum tocopherols. Concentrations of various constituents in blood of dairy cows during stages of terminal gestation and initial lactation. I., 503.
— self-selected, riboflavin metabolism of young women on, 435.
— similar to those consumed by certain low income groups in South Carolina, use of enriched and non-enriched flour in, 421.
Dietary regime, normal, effect of fat on calcium and phosphorus metabolism in normal growing rats under, 165.
Differences in influence of live yeast on absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by human subjects, and effect of distribution of vitiamin doses, 225.
Digestibility of rapeseed and cottonseed oils in human subjects, 369.
Dogs fed protein hydrolysates or amino acids, urinary excretion of amino acids and peptides by, 155.
— protein-depleted, effect of retention of nitrogen in casein or lactalbumin hydrolysates on regeneration of plasma proteins of, 319.

EGG, biological values relative to, and replacement values of some cereal proteins in human subjects, 405.
— embryonated hen, distribution of thiamine in. I., 237; II., 267.
ELLIS, N. R. See Hoagland, R., 381.
ELVEHJEM, C. A. See Couch, J. R., 105; Ramasarma, G. B., 177; Tove, S. B., 469.
Embryonated hen egg, distribution of thiamine in. I., 237; II., 267.
Embryonic tissues, content of. Distribution of thiamine in embryonated hen egg. II., 267.
Endemic beriberi (see Beriberi, endemic).
Energy expenditure of boys and girls 9 to 11 years of age: (1) sitting listening to the radio (phonograph), (2) sitting singing, and (3) standing singing, 1.
Enhancement of nutritive value of wheat gluten by supplementation with lysine, 331.
Enriched flour (see Flour, enriched).
Enrichment, artificial, of white rice as solution to endemic beriberi: I. Preliminary report of field trials, 443.
EVANS, H. M. See Nelson, M. M., 11.
Excretion of amino acids by rats, and their growth, effect of fat in diet on, 257.
— urinary (see Urinary excretion).

FAT(S), comparative nutritive value of. XI., 361; XII., 369.
— effect on calcium and phosphorus metabolism in normal growing rats under normal dietary regime, 165.
— in diet of rats, effect on growth and excretion of amino acids, 257.
— influence of high levels of, with suboptimum levels of riboflavin, on growth of chicks, 247.
Fertilization, ovulation, and transport of ova in old, vitamin E deficient rats, 97.
Field trials, preliminary report of. Artificial enrichment of white rice as solution to endemic beriberi: I., 443.
FITZPATRICK, W. H. See Swain, H. L., 63.
Flour, enriched and non-enriched, use in diet similar to those consumed by certain low income groups in South Carolina, 421.
— improving the nutritive value of. III., 421.
Folic acid, microbiological assay of, as applied to milk, 25.
— — studies in mink and fox, 469
FOLKVORD, S. See Kringstad, H., 489.
Food consumption in various U. S. Army camps, simultaneous surveys of, 51.
— — of soldiers in subarctic climate (Fort Churchill, Manitoba, Canada, 1947–48), 63.
Foodstuffs, several uncooked and cooked Mexican, determination of some essential amino acids in, 293.
Fowl, mature, utilization of lactose by, 105.
Fox, folic acid studies in, 469.
FRENCHMAN, R. See Johnston, F. A., 479.
FUKUI, T. See Deuel, H. J. Jr., 361.

GARBER, M., M. M. MARQUETTE AND H. T. PARSONS. The availability of vitamins from yeasts. V. Differences in the influence of live yeast on the absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by human subjects, and the effect of distribution of vitamin doses. 225.
GARDNER, J. See Deuel, H. J., Jr., 369.

Gestation, terminal, and initial lactation, concentrations of various constituents in blood of dairy cows during stages of, I., 503.
Girls (see Children).
Gluten, wheat (see Wheat gluten).
Goitrogenic properties of soybeans, relation of iodine to, 115.
GOODING, C. M. See Deuel, H. J., Jr., 361.
GOYCO, J. A., AND C. F. ASENJO. Net protein and growth-promoting values of three different types of yeast prepared under identical conditions, 517.
GREENBERG, S. M. See Deuel, H. J., Jr., 361.
Growth of chicks, influence of high levels of fat with suboptimum levels of riboflavin on, 247.
— of rats and their excretion of amino acids, effect of fat in diet on, 257.
Growth-promoting activity, possible of Δ^{12}-octadecenoic acid, 361.
— and net protein values of three different types of yeast prepared under identical conditions, 517.
Guinea pigs, wrist stiffness syndrome in, 87.
GUZMÁN, J. See Massieu, G. H., 293.

HALVERSON, A. W., M. ZEPPLIN AND E. B. HART. Relation of iodine to the goitrogenic properties of soybeans, 115.
Hamster, Syrian, vitamin B_6 deficiency in, 131.
HANKINS, O. G. See Hoagland, R., 381.
HARRIS, J. N., AND F. I. SCOULAR. Riboflavin metabolism of young women on self-selected diets, 435.
HART, E. B. See Halverson, A. W., 115.
HEIMER, C. B., H. L. MASLOW AND A. E. SOBEL. Influence of thyroid on utilization of vitamin A, 345.
Hen egg (see Egg, hen).
HENDERSON, L. M. See Ramasarma, G. B., 177.
HOAGLAND, R., O. G. HANKINS, N. R. ELLIS AND G. G. SNIDER. Nutritive properties of protein in different cuts of beef, 381.
HODSON, A. Z. The microbiological assay of "folic acid" as applied to milk, 25.
HOFFMAN, W. S., AND G. C. MCNEIL. The enhancement of the nutritive value of wheat gluten by supplementation with lysine, as determined from nitrogen balance indices in human subjects, 331.
HUGHES, J. S. See Latschar, C. E., 503.
Human subjects (see Man).
Hydrochloride (pure thiamine), pure riboflavin and nitrogen, differences in influence of live yeast on absorption by human subjects, and effect of distribution of vitamin doses, 225.
Hydrolysates (casein or lactalbumin), effect of retention of nitrogen in, on regeneration of plasma proteins of protein-depleted dogs, 319.
— (protein) or amino acids, urinary excretion of amino acids and peptides by dogs fed, 155.

INFANT at birth, relation of maternal nutrition to condition of: study of siblings, 453.
— male, biological value of corn and wheat proteins in, with note on utilization of D-tryptophan, 215.
INSTITUTE OF NUTRITION (see American Institute of Nutrition).
Iodine, relation to goitrogenic properties of soybeans, 115.
Iron metabolism (see Metabolism, iron).

JAMES, M. F. See Neumann, A. L., 195.
JOHNSON, B. C. See Neumann, A. L., 195.
JOHNSON, R. M. See Deuel, H. J. Jr., 369.
JOHNSTON, F. A., R. FRENCHMAN AND E. D. BOROUGHS. The iron metabolism of young women on two levels of intake, 479.

KAUNITZ, H. See Blandau, R. J., 97.
KENNEDY, C. See Calverley, C. E., 165.
KEMMERER, A. R., AND R. ACOSTA. The essential amino acid content of several vegetables, 527.
KIRK, E., AND M. CHIEFFI. Vitamin studies in middle-aged and old individuals. III. Thiamine and pyruvic acid blood concentrations, 353.
KOEHN, C. J. See Swain, H. L., 63.
KRAUSE, R. F. A paradoxical relationship between serum level and liver content of vitamin A, 535.
KRIDER, J. L. See Neumann, A. L., 195.
KRINGSTAD, H., AND S. FOLKVORD. The nutritive value of cod roe and cod liver, 489.

LACTALBUMIN or casein hydrolysates, effect of retention of nitrogen in, on regeneration of plasma proteins of protein-depleted dogs, 319.
Lactation, initial, and terminal gestation, concentrations of various constituents in blood of dairy cows during stages of, I., 503.
Lactose, utilization by mature fowl, 105.
LATSCHAR, C. E., G. H. WISE, D. B. PARRISH AND J. S. HUGHES. Concentrations of various constituents in blood of dairy cows during stages of terminal gestation and initial lactation. I. Effect of prepartal diet on serum tocopherols, 503.
LEDUC, E. H. See Wilson, J. W., 73.
LEIN, M. See Albanese, A. A., 215.
Lesions, carious, in cotton rat molars. III., 275.
LINN, D. R. See Westerman, B. D., 421.
Live yeast (see Yeast, live).
Liver (cod) and cod roe, nutritive value of, 489.
— content and serum level of vitamin A, paradoxical relationship between, 535.
Low income groups in South Carolina, use of enriched and non-enriched flour in diets similar to those consumed by certain, 421.
Lysine, enhancement of nutritive value of wheat gluten by supplementation with, 331.

MALE infant, biological value of corn and wheat proteins in, with note on utilization of D-tryptophan, 215.
Man, differences in influence of live yeast on absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by, and effect of distribution of vitamin doses, 225.
— digestibility of rapeseed and cottonseed oils in, 369.
— enhancement of nutritive value of wheat gluten by supplementation with lysine, as determined from nitrogen balance indices in, 331.
— relative to egg biological values and replacement values of some cereal proteins in, 405.
MARQUETTE, M. M. See Garber, M., 225.
MASLOW, H. L. See Heimer, C. B., 345.

Massieu, G. H., J. Guzmán, R. O. Cravioto and J. Calvo. Determination of some essential amino acids in several uncooked and cooked Mexican foodstuffs, 293.
Maternal nutrition, relation to condition of infant at birth: study of siblings, 453.
Mature fowl (see Fowl, mature).
Maynard, L. A. See Smith, S. E., 87.
McNeil, G. C. See Hoffman, W. S., 331.
Mead Johnson and Company B complex award, 1949, presentation, 402.
Metabolism (calcium and phosphorus) in normal growing rats under normal dietary regime, effect of fat on, 165.
— effects of thiouracil on, 41.
— iron, of young women on two levels of intake, 479.
— riboflavin, of young women on self-selected diets, 435.
Mexican foodstuffs, several uncooked and cooked, determination of some essential amino acids in, 293.
Microbiological assay of "folic acid" as applied to milk, 25.
Middle-aged individuals, vitamin studies in. III., 353.
Milk, microbiological assay of "folic acid" as applied to, 25.
Mink, folic acid studies in, 469.
Molars (cotton rat), carious lesions in. III., 275.
Mouse, production of biotin deficiency in, 73.
Murlin, J. R. See Clark, L. C., 405.

Nelson, M. M., and H. M. Evans. Pteroylglutamic acid and reproduction in the rat, 11.
Net protein and growth-promoting values of three different types of yeast prepared under identical conditions, 517.
Neumann, A. L., J. L. Krider, M. F. James and B. C. Johnson. The choline requirement of the baby pig, 195.
Nitrogen balance indices in human subjects, enhancement of nutritive value of wheat gluten by supplementation with lysine, as determined from, 331
— effect of retention, in casein or lactalbumin hydrolysates on regeneration of plasma proteins of protein-depleted dogs, 319
— for growing rat, purified amino acids as source of, 177
— pure thiamine hydrochloride and pure riboflavin, differences in influence of live yeast on absorption by human subjects, and effect of distribution of vitamin doses, 225
Non-enriched flour (see Flour, enriched and non-enriched)
Nutrition, American Institute of (see American Institute of Nutrition)
— Borden award in (see Borden award).
— (maternal), relation to condition of infant at birth: study of siblings, 453.
— studies during pregnancy. V. Relation of maternal nutrition to condition of infant at birth: study of siblings, 453.
Nutritive properties of protein in different cuts of beef, 381.
— value of cod roe and cod liver, 489.
— — of fats, comparative. XI. Possible growth-promoting activity of Δ^{13}-octadecenoic acid, 361; XII. Digestibility of rapeseed and cottonseed oils in human subjects, 369.
— — of flour, improving the. III. Use of enriched and non-enriched flour in diets similar to those consumed by certain low income groups in South Carolina, 421.
— — of wheat gluten, enhancement by supplementation with lysine, 331.

Octadecenoic acid (Δ^{13}-), possible growth-promoting activity of, 361.
Old individuals, vitamin studies in. III., 353.
— vitamin E deficient rats, ovulation, fertilization and transport of ova in, 97.
Osborne and Mendel award, presentation, 1949, 402.
Ovulation, fertilization, and transport of ova in old, vitamin E deficient rats, 97.

Panganiban, E. C. See Salcedo, J., Jr., 443.
Panzer, F. See Pearson, P. B., 257.
Paradoxical relationship between serum level and liver content of vitamin A, 535.
Parrish, D. B. See Latschar, C. E., 503.
Parsons, H. T. See Garber, M., 225.
Particle size and consistency of purified ration, effect on incidence and type of carious lesions in cotton rat molars, 275.
Pearson, P. B., and F. Panzer. Effect of fat in the diet of rats on their growth and their excretion of amino acids, 257.
— See Reiser, R., 247.
Pedroche, A. See Salcedo, J., Jr., 443.
Peptides and amino acids, urinary excretion by dogs fed protein hydrolysates or amino acids, 155.
Phonograph (radio), energy expenditure of boys and girls 9 to 11 years of age sitting listening to, 1.
Phosphorus metabolism (see Metabolism, phosphorus).
Pig, baby, choline requirement of, 195.
— guinea (see Guinea pig).
Plasma proteins of protein-depleted dogs, effect of retention of nitrogen in casein or lactalbumin hydrolysates on regeneration of, 319.
Porter, C. C. See Silber, R. H., 155.
Porter, W. E. See Scrimshaw, N. S., 237, 267.
Pregnancy, nutrition studies during. V., 453.
Prepartal diet, effect on serum tocopherols. Concentrations of various constituents in blood of dairy cows during stages of terminal gestation and initial lactation. I., 503.
Protein(s) (corn and wheat), biological value in male infant, with note on utilization of D-tryptophan, 215.
— depleted dogs, effect of retention of nitrogen in casein or lactalbumin hydrolysates on regeneration of plasma proteins of, 319.
— hydrolysates or amino acids, urinary excretion of amino acids and peptides by dogs fed, 155.
— in different cuts of beef, nutritive properties of, 381.
— (net) and growth-promoting values of three different types of yeast prepared under identical conditions, 517.
— (plasma) of protein-depleted dogs, effect of retention of nitrogen in casein or lactalbumin hydrolysates on regenreration of, 319.
— (some cereal) in human subjects, relative to egg biological values and replacement values of, 405.
Pteroylglutamic acid and reproduction in rat, 11.
Purified amino acids as source of nitrogen for growing rat, 177.
— ration, consistency of, and particle size, effect on incidence and type of carious lesions in cotton rat molars, 275.

Pyk, O. F. See Taylor, C. M., 1.
Pyruvic acid and thiamine blood concentrations. Vitamin studies in middle-aged and old individuals. III., 353.

QUERCETIN and rutin, value in scurvy, 305.

RADIO (phonograph), energy expenditure of boys and girls 9 to 11 years of age sitting listening to, 1.
Ramasarma, G. B., L. M. Henderson and C. A. Elvehjem. Purified amino acids as a source of nitrogen for the growing rat, 177.
Rapeseed and cottonseed oils, digestibility in human subjects, 369.
Rat(s) (cotton) molars, carious lesions in. III., 275.
— effect of fat in diet of, on growth and excretion of amino acids, 257.
— growing, purified amino acids as source of nitrogen for, 177.
— normal growing, under normal dietary regime, effect of fat on calcium and phosphorus metabolism in, 165.
— (old, vitamin E deficient), ovulation, fertilization and transport of ova in, 97.
— pteroylglutamic acid and reproduction in, 11.
Ration, consistency of purified, and particle size, effect on incidence and type of carious lesions in cotton rat molars, 275.
Regime, dietary (see Dietary regime).
Reiser, R., and P. B. Pearson. The influence of high levels of fat with suboptimum levels of riboflavin on the growth of chicks, 247.
Replacement values and relative to egg biological values of some cereal proteins in human subjects, 405.
Reproduction and pteroylglutamic acid in rat, 11.
Riboflavin, influence on growth of chicks of high levels of fat with suboptimum levels of, 247
— metabolism (see Metabolism, riboflavin).
— (pure), pure thiamine hydrochloride and nitrogen, differences in influence of live yeast on absorption by human subjects, and effect of distribution of vitamin doses, 225.
Rice, white, artificial enrichment of, as solution to endemic beriberi. I. Preliminary report of field trials, 443.
Roe, cod (see Cod roe).
Rutin and quercetin, value in scurvy, 305.
Ryer, H. See Clark, L. C., 405.

SALCEDO, J., Jr., A. Pedroche, E. C. Panganiban and J. F. de Leon. Artificial enrichment of white rice as a solution to endemic beriberi. I. Preliminary report of field trials, 443.
Schaefer, A. E. See Tove, S. B., 469.
Schor, H. C., and H. L. Swain. Simultaneous surveys of food consumption in various camps of the U. S. Army, 51.
Scoular, F. I. See Harris, J. N., 435.
Scrimshaw, M. W. See Scrimshaw, N. S., 237, 267.
Scrimshaw, N. S., W. E. Porter and M. W. Scrimshaw. The distribution of thiamine in the embryonated hen egg. I. The content of the whole embryo, 237; II. The content of embryonic tissues, 267.
Scurvy, value of rutin and quercetin in, 305.
Self-selected diets, riboflavin metabolism of young women on, 435.
Serum level and liver content of vitamin A, paradoxical relationship between, 535.
— tocopherols, effect of prepartal diet on. Concentrations of various constituents in blood of dairy cows during stages of terminal gestation and initial lactation. I., 503.
Shaw, J. H. Carious lesions in cotton rat molars. III. Effect of the particle size and consistency of the purified ration on the incidence and type of carious lesions, 275.
Shwartzman, G., and L. Strauss. Vitamin B_6 deficiency in the Syrian hamster, 131.
Siblings, study of: Relation of maternal nutrition to condition of infant at birth, 453.
Silber, R. H., and C. C. Porter. Urinary excretion of amino acids and peptides by dogs fed protein hydrolysates or amino acids, 155.
Simultaneous surveys of food consumption in various U. S. Army camps, 51.
Singing, energy expenditure of boys and girls 9 to 11 years of age sitting and standing, 1.
Slanetz, C. A. See Blandau, R. J., 97.
Smetak, E. M. See Albanese, A. A., 215.
Smith, S. E., M. A. Williams, A. C. Bauer and L. A. Maynard. The wrist stiffness syndrome in guinea pigs, 87.
Snider, G. G. See Hoagland, R., 381.
Snyderman, S. E. See Albanese, A. A., 215.
Sobel, A. E. See Heimer, C. B., 345.
Soldiers, food consumption in subarctic climate (Fort Churchill, Manitoba, Canada, 1947–48), 63.
Sostman, E. R. See Taylor, C. M., 1.
Soybeans, relation of iodine to goitrogenic properties of, 115.
Stevenson, S. S. See Burke, B. S., 453.
Stiffness (wrist) syndrome in guinea pigs, 87.
Straub, E. E. See Deuel, H. J., Jr., 361.
Strauss, L. See Shwartzman, G., 131.
Stuart, H. C. See Burke, B. S., 453.
Subarctic climate, food consumption of soldiers in (Fort Churchill, Manitoba, Canada, 1947–48), 63.
Sunde, M. L. See Couch, J. R., 105.
Surveys, simultaneous, of food consumption in various U. S. Army camps, 51.
Swain, H. L., F. M. Toth, F. C. Consolazio, W. H. Fitzpatrick, D. I. Allen and C. J. Koehn. Food consumption of soldiers in a subarctic climate (Fort Churchill, Manitoba, Canada, 1947–48), 63.
— See Schor, H. C., 51.
Swift, R. W. See Bratzler, J. W., 41.
Syrian hamster, vitamin B_6 deficiency in, 131.

TAYLOR, C. M., O. F. Pyke, A. B. Caldwell and E. R. Sostman. The energy expenditure of boys and girls 9 to 11 years of age: (1) sitting listening to the radio (phonograph), (2) sitting singing, and (3) standing singing, 1.
Templeton, F. See Westerman, B. D., 421.
Thiamine and pyruvic acid blood concentrations. Vitamin studies in middle-aged and old individuals. III., 353.
— distribution in embryonated hen egg. I. Content of whole embryo, 237; II. Content of embryonic tissues, 267.
— hydrochloride (pure), pure riboflavin and nitrogen, differences in influence of live yeast on absorption by human subjects, and effect of distribution of vitamin doses, 225.
Thiouracil, effects on metabolism, 41.
Thomas, B. See Deuel, H. J. Jr., 369.
Thyroid, influence on utilization of vitamin A, 345.

Tissues, content of embryonic. Distribution of thiamine in embryonated hen egg. II. 267.
Tocopherols, serum, effect of prepartal diet on. Concentrations of various constituents in blood of dairy cows during stages of terminal gestation and initial lactation. I., 503.
TOTH, F. M. See Swain, H. L., 63.
TOVE, S. B., A. E. SCHAEFER AND C. A. ELVEHJEM. Folic acid studies in the mink and fox, 469.
Tryptophan (D-), biological value of corn and wheat proteins in male infant, with note on utilization of, 215.

UNITED States Army, simultaneous surveys of food consumption in various camps of, 51.
Urinary excretion of amino acids and peptides by dogs fed protein hydrolysates or amino acids, 155.

VEGETABLES, essential amino acid content of several, 527.
VESTAL, B. See Albanese, A. A., 215.
Vitamin(s) A, influence of thyroid on utilization of, 345.
— — paradoxical relationship between serum level and liver content of, 535.
— availability from yeasts. V. Differences in influence of live yeast on absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by human subjects, and effect of distribution of vitamin doses, 225.
— B_6 deficiency in Syrian hamster, 131.
— doses, differences in influence of live yeast on absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by human subjects, and effect of distribution of, 225.
— E deficient, old rats, ovulation, fertilization and transport of ova in, 97.
— studies in middle-aged and old individuals. III. Thiamine and pyruvic acid blood concentrations, 353.

WELLS, R. I. See Westerman, B. D., 421.
WESTERMAN, B. D., D. R. LINN, F. TEMPLETON AND R. I. WELLS Improving the nutritive value of flour. III. The use of enriched and non-enriched flour in diets similar to those consumed by certain low income groups in South Carolina, 421.
Wheat and corn proteins, biological value in male infant, with note on utilization of D-tryptophan, 215.
— gluten, enhancement of nutritive value of, by supplementation with lysine, 331.
WILLIAMS, M. A. See Smith, S. E., 87.
WILSON, J. W., E. H. LEDUC AND D. H. WINSTON. The production of biotin deficiency in the mouse, 73.
WINSTON, D. H. See Wilson, J. W., 73.
WISE, G. H. See Latschar, C. E., 503.
Women, young, iron metabolism of, on two levels of intake, 479.
— — on self-selected diets, riboflavin metabolism of, 435.
WORCESTER, J. See Burke, B. S., 453.
Wrist stiffness syndrome in guinea pigs, 87.

YEAST(S), availability of vitamins from. V., 225.
— (live), differences in influence of, on absorption of pure thiamine hydrochloride, pure riboflavin and nitrogen by human subjects, and effect of distribution of vitamin doses, 225.
— three different types of, prepared under identical conditions, net protein and growth-promoting values of, 517.
Young women (see Women, young).

ZEPPLIN, M. See Halverson, A. W., 115.